

Page 1 of 16 05/31/2011 Percent of Surgery Patients given an antibiotic at the right time (within one hour before surgery) to help prevent infection

Page 2 of 16 05/31/2011

Number of Patients-2	Footnote-2	

Page 3 of 16 05/31/2011 Percent of Surgery Patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)

Page 4 of 16 05/31/2011

Number of Patients-3	Footnote-3

Page 5 of 16 05/31/2011

Percent of Surgery Patients who were given the right kind of antibiotic to help prevent infection

> Page 6 of 16 05/31/2011

Number of Patients-4	Footnote-4

Page 7 of 16 05/31/2011



Percent of Surgery Patients who got treatment at right time (within 24 hours before or after surgery) to help prevent blood clot

Page 8 of 16 05/31/2011

Number of Patients-5	Footnote-5

Page 9 of 16 05/31/2011



Percent of Surgery Patients whose doctors ordered treatments to prevent blood clots after certain types of surgeries

Page 10 of 16 05/31/2011

Number of Patients-6	Footnote-6

Page 11 of 16 05/31/2011 Percent of all Heart Surgery Patients whose blood sugar is kept under good control in the days right after surgery

Page 12 of 16 05/31/2011

Number of Patients-7	Footnote-7

Page 13 of 16 05/31/2011 Percent of Surgery Patients needing hair removed from the surgical area before surgery who had hair removed using a safer method

Page 14 of 16 05/31/2011

Number of Patients-8	Footnote-8	Surgery patients who were taking heart drugs called beta blockers before coming to the hospital, who were kept on them
----------------------	------------	--

Page 15 of 16 05/31/2011

Outpatients having surgery who got an antibiotic at the right time within one hour before surgery (higher numbers are better)

Outpatients having surgery who got the right kind of antibiotic (higher numbers are better)

> Page 16 of 16 05/31/2011